
Trigonometric Equations and Identities - Edexcel Past Exam Questions

1. (a) Show that the equation

$$5 \cos^2 x = 3(1 + \sin x)$$

can be written as

$$5 \sin^2 x + 3 \sin x - 2 = 0. \quad (2)$$

- (b) Hence solve, for $0 \leq x < 360^\circ$, the equation

$$5 \cos^2 x = 3(1 + \sin x),$$

giving your answers to 1 decimal place where appropriate. (5)

Jan 05 Q4

2. Solve, for $0 \leq x \leq 180^\circ$, the equation

(a) $\sin(x + 10^\circ) = \frac{\sqrt{3}}{2}, \quad (4)$

(b) $\cos 2x = -0.9$, giving your answers to 1 decimal place. (4)

June 05 Q5

3. (a) Find all the values of θ , to 1 decimal place, in the interval $0^\circ \leq \theta < 360^\circ$ for which

$$5 \sin(\theta + 30^\circ) = 3. \quad (4)$$

- (b) Find all the values of θ , to 1 decimal place, in the interval $0^\circ \leq \theta < 360^\circ$ for which

$$\tan^2 \theta = 4. \quad (5)$$

Jan 06 Q8



-
4. (a) Given that $\sin \theta = 5 \cos \theta$, find the value of $\tan \theta$. (1)

(b) Hence, or otherwise, find the values of θ in the interval $0 \leq \theta < 360^\circ$ for which

$$\sin \theta = 5 \cos \theta,$$

giving your answers to 1 decimal place. (3)

June 06 Q6

5. (a) Show that the equation

$$3 \sin^2 \theta - 2 \cos^2 \theta = 1$$

can be written as

$$5 \sin^2 \theta = 3. \quad (2)$$

(b) Hence solve, for $0^\circ \leq \theta < 360^\circ$, the equation

$$3 \sin^2 \theta - 2 \cos^2 \theta = 1,$$

giving your answer to 1 decimal place. (7)

Jan 08 Q4

6. Solve, for $0 \leq x < 360^\circ$,

(a) $\sin(x - 20^\circ) = \frac{1}{\sqrt{2}}$, (4)

(b) $\cos 3x = -\frac{1}{2}$. (6)

June 08 Q9

7. (a) Show that the equation

$$4 \sin^2 x + 9 \cos x - 6 = 0$$

can be written as

$$4 \cos^2 x - 9 \cos x + 2 = 0. \quad (2)$$

(b) Hence solve, for $0 \leq x < 720^\circ$,

$$4 \sin^2 x + 9 \cos x - 6 = 0,$$

giving your answers to 1 decimal place. (6)

Jan 09 Q8



8. (i) Solve, for
- $-180^\circ \leq \theta < 180^\circ$
- ,

$$(1 + \tan \theta)(5 \sin \theta - 2) = 0. \quad (4)$$

- (ii) Solve, for
- $0 \leq x < 360^\circ$
- ,

$$4 \sin x = 3 \tan x. \quad (6)$$

June 09 Q7

9. (a) Show that the equation

$$5 \sin x = 1 + 2 \cos^2 x$$

can be written in the form

$$2 \sin^2 x + 5 \sin x - 3 = 0. \quad (2)$$

- (b) Solve, for
- $0 \leq x < 360^\circ$
- ,

$$2 \sin^2 x + 5 \sin x - 3 = 0. \quad (4)$$

Jan 10 Q2

10. (a) Given that
- $5 \sin \theta = 2 \cos \theta$
- , find the value of
- $\tan \theta$
- .

(1)

- (b) Solve, for
- $0 \leq x < 360^\circ$
- ,

$$5 \sin 2x = 2 \cos 2x,$$

giving your answers to 1 decimal place.

(5)**June 10 Q5**



11. (a) Show that the equation

$$3 \sin^2 x + 7 \sin x = \cos^2 x - 4$$

can be written in the form

$$4 \sin^2 x + 7 \sin x + 3 = 0. \quad (2)$$

- (b) Hence solve, for $0 \leq x < 360^\circ$,

$$3 \sin^2 x + 7 \sin x = \cos^2 x - 4$$

giving your answers to 1 decimal place where appropriate.

(5)

Jan 11 Q7

12. Solve for $0 \leq x < 360^\circ$, giving your answers in degrees to 1 decimal place,

$$3 \sin (x + 45^\circ) = 2. \quad (4)$$

June 11 Q7(*edited*)
